



Optimizing Atrazine Application Rates for Efficacious Weed Control in Maize Cultivation

Muneeba¹, Faran Muhammad¹, Rashid Khan², Muhammad Shafique³

¹Department of Agronomy, University of Agriculture Faisalabad

²Horticulture section, Agriculture Research Institute Dera Ismail Khan

³Sugarcane Research institute, Ayub Research Faisalabad

ARTICLE INFO

Article History:

Received:	February	15, 2024
Revised:	March	20, 2024
Accepted:	April	15, 2024
Available Online:	May	10, 2024

Keywords:

experimental design incorporates

ABSTRACT

This study delves into the intricate task of optimizing atrazine application rates to achieve efficacious weed control in maize cultivation. Atrazine, a widely employed herbicide known for its effectiveness against various weed species, is a cornerstone in contemporary weed management strategies. The challenge lies in identifying the precise application rates that strike a balance between robust weed eradication and minimizing potential ecological impacts. Through meticulous field trials and systematic data analyses, this research systematically explores a range of atrazine application rates to discern their differential effects on weed populations, crop health, and overall maize productivity. The experimental design incorporates varying concentrations (0, 0.5, 1, 1.5 and 2 ml L⁻¹) of atrazine, allowing for a comprehensive evaluation of its impact on both target weeds and the maize crop. Parameters such as weed density, species composition, crop vigor, and yield components are rigorously assessed. The study aims to elucidate the optimal atrazine application rates that maximize weed control efficacy while minimizing the risk of adverse effects on non-target organisms and environmental sustainability. The anticipated outcomes of this research hold significant implications for sustainable agriculture, providing practitioners with data-driven insights to refine atrazine application practices. By offering a nuanced understanding of the intricate relationship between atrazine dosages and weed control outcomes, this study contributes to the ongoing discourse on precision herbicide application in maize cultivation. Ultimately, the findings aim to guide farmers, agronomists, and policymakers towards more informed and sustainable weed management practices in maize crops.



© 2024 The Authors, Published by AIRSD. This is an Open Access Article under the Creative Common Attribution Non-Commercial 4.0

Corresponding Author's Email: faran0169@gmail.com

INTRODUCTION

The intricate dance between crop productivity and weed interference stands as a perennial challenge in modern agriculture ¹. Among the myriad tools at the disposal of farmers, herbicides play a pivotal role in shaping the delicate equilibrium between maximizing yields and mitigating the detrimental impacts of weed competition ². Atrazine, a triazine-class herbicide, has long been a cornerstone in the arsenal of weed management strategies, particularly in maize cultivation ³. Its broad-spectrum effectiveness and versatility make it a go-to solution for controlling a spectrum of weed species that encroach upon the vitality of maize crops ⁴.

While atrazine's efficacy is undisputed, the optimization of application rates represents a critical juncture in precision agriculture ⁵. Striking the delicate balance between effective weed controls and minimizing potential ecological repercussions necessitates a nuanced understanding of atrazine's dose-response relationship ⁶. This research embarks on a journey to unravel this complexity, seeking to optimize atrazine application rates for efficacious weed control in maize cultivation.

Maize, or corn (*Zea mays*), stands as one of the world's primary staple crops, sustaining both human and livestock populations. Its significance in global food security underscores the imperative of optimizing cultivation practices to ensure robust yields. Weeds, however, pose a perennial threat, competing for essential resources and hampering the growth and productivity of maize crops. In this context, herbicides have emerged as indispensable tools, offering a targeted and efficient means of weed control ^{7,8}.

Atrazine, a chlorotriazine herbicide, has been a linchpin in the realm of weed management for decades. Its mode of action involves inhibiting photosynthesis in susceptible plants, rendering it effective against a broad spectrum of grasses and broadleaf weeds. Its residual activity further extends its effectiveness, providing a prolonged shield against weed resurgence ⁹. Despite its efficacy, the environmental impact of atrazine has sparked debates, necessitating a nuanced approach to its application.

The optimization of atrazine application rates becomes particularly crucial for several reasons. First, excessive application may lead to environmental contamination, affecting non-target plants and organisms, and potentially leaching into water sources. Second, economic considerations prompt the need for judicious herbicide use, ensuring cost-effectiveness for farmers while maintaining efficacy. Third, evolving weed populations may exhibit varying degrees of susceptibility, demanding a tailored approach to dosage.

The substantial significance for agricultural practitioners, researchers, and policymakers alike. By unraveling the intricate relationship between atrazine application rates and weed control efficacy, the study contributes to the development of more sustainable and precise weed management practices in maize cultivation. The findings hold the potential to inform agronomic decisions, guiding farmers towards optimized herbicide use that aligns with both economic and environmental considerations.

As agriculture navigates the complex terrain of feeding a growing global population while minimizing environmental impacts, the optimization of herbicide application rates emerges as a crucial strategy.

This research endeavors to systematically assess the impact of varying atrazine application rates on weed populations (weed density and species composition) and growth and yield of maize.

This study, focused on atrazine in maize cultivation, aims to carve a path towards a more nuanced, efficient, and sustainable approach to weed management, thereby contributing to the broader discourse on precision agriculture and responsible herbicide use.

MATERIALS AND METHODS

Experimental Design:

The study employed a randomized complete block design (RCBD) to account for potential spatial variability in the experimental field. A total of six treatment levels were established, representing atrazine application rates of 0, 0.5, 1, 1.5, and 2 ml per liter of herbicide solution.

Field Site Selection:

A representative maize cultivation site was selected based on uniform soil characteristics and historical weed management practices. The site had not been subjected to recent herbicide applications to avoid residual effects.

Herbicide Application:

Atrazine, a chlorotriazine herbicide, was used as the primary weed control agent. A range of application rates, including 0, 0.5, 1, 1.5 and 2 ml per liter, were prepared to encompass a spectrum of dosage levels. Herbicide application was carried out during the early stages of maize growth, corresponding to the recommended timing for effective weed control.

Plot Preparation:

Experimental plots were demarcated with suitable spacing to prevent herbicide drift and facilitate proper replication. Each treatment level was replicated across multiple plots to ensure robust statistical analyses.

Weed Density Assessment:

Weed density was assessed by systematically sampling a predetermined area within each plot. Weed species, density, and diversity were recorded to evaluate the herbicide's efficacy against different weed types.

Maize Growth Parameters:

Maize growth parameters, including plant height and chlorophyll content, were measured at regular intervals throughout the growing season. Plant height provided insights into crop vigor, while chlorophyll content served as an indicator of overall plant health.

Grain Yield Measurement:

Maize grain yield was determined by harvesting mature maize cobs from each plot. Harvested grain was thoroughly cleaned, weighed, and expressed on a per-hectare basis for standardized comparison.

Biomass and Straw Yield:

Above-ground biomass, comprising both grain and vegetative plant components, was collected from each plot at the time of harvest. Separation of grain and straw components facilitated the quantification of biomass and straw yield.

Statistical Analysis:

Statistical analyses, including analysis of variance (ANOVA), were employed to discern significant differences among the atrazine application rates for each parameter. Post-hoc tests were conducted where necessary to identify specific treatment effects.

Replicability and Randomization:

The experimental design incorporated a sufficient number of replications for each treatment to enhance statistical power. Randomization of treatment application and data collection points minimized bias and increased the robustness of the study.

Data Recording and Documentation:

All experimental procedures, including herbicide preparation, application, and data collection, were meticulously recorded. The documentation included dates, weather conditions, and any unforeseen events that could influence study outcomes.

Safety Precautions:

Adherence to safety protocols during herbicide handling and application was paramount to minimize risks to researchers, the environment, and neighboring ecosystems. Herbicide containers and waste were disposed of in accordance with environmental safety guidelines.

Environmental Monitoring:

Throughout the study, environmental conditions such as soil moisture, temperature, and weather patterns were monitored. These variables were considered in data interpretation to contextualize the herbicide's effects on both weed and crop responses.

RESULTS

Weed Population Dynamics:

Atrazine application exhibited a dose-dependent response in weed density. The control group (0 ml/L) had the highest weed density, while increasing atrazine rates correlated with a significant reduction in weed populations. At 2 ml/L, weed density reached its lowest point, indicating the efficacy of atrazine in suppressing weed growth. Atrazine demonstrated selectivity in weed control, influencing different weed species to varying extents. Broadleaf weeds showed higher susceptibility to atrazine, with a noticeable decline in their representation as herbicide rates increased. Grass species also exhibited reduced density with higher atrazine doses, highlighting the herbicide's effectiveness against both weed categories (Figure 1).




Figure 1. Effect of different rates of atrazine on number and diversity of weeds in maize crop

Plant Height and Chlorophyll:

The inverse relationship between atrazine application rates and weed density aligns with the herbicide's mode of action, inhibiting photosynthesis and impeding weed growth. The selectivity observed in weed species composition emphasizes atrazine's differential impact on broadleaf and grassy weeds. Such selectivity is crucial in maintaining crop integrity while efficiently managing weed populations. The positive correlation between atrazine application rates and maize grain yield underscores the herbicide's pivotal role in optimizing crop productivity. Higher atrazine doses effectively reduced weed competition, allowing maize plants to allocate resources more efficiently toward grain production (Figure 2).




Figure 2. Effect of different herbicides on plant height of maize crop

Grain Yield:

Maize grain yield demonstrated a clear positive correlation with atrazine application rates. The control group exhibited the lowest grain yield, while incremental increases in atrazine dosage resulted in a significant improvement in maize productivity. At 1.5 ml/L, maize grain yield

reached its peak, emphasizing the potential for optimizing atrazine application rates to maximize crop output (Figure 3).





Figure 3. Effect of different herbicides on grain yield of maize crop

Biomass Yield and Straw Yield:

Atrazine application influenced both biomass and straw yield, reflecting the herbicide's impact on overall maize development. Biomass yield increased steadily with atrazine dosage, indicating robust crop growth and effective weed suppression. Straw yield followed a similar trend, demonstrating the herbicide's ability to enhance not only grain yield but also vegetative plant components. (Figure 4 a, b).






Figure 4. Effect of different rates of atrazine on straw (a) and biomass (b) yield of maize crop

DISCUSSION

The inverse relationship between atrazine application rates and weed density aligns with the herbicide's mode of action, inhibiting photosynthesis and impeding weed growth. The selectivity observed in weed species composition emphasizes atrazine's differential impact on broadleaf and grassy weeds¹⁰. Such selectivity is crucial in maintaining crop integrity while efficiently managing weed populations.

The observed reduction in maize plant height with increasing atrazine application rates aligns with the herbicide's mode of action, which inhibits photosynthesis in susceptible plants. The dose-dependent phytotoxic effect on plant height underscores the importance of careful consideration when optimizing atrazine application rates to balance weed control efficacy with potential impacts on crop development¹⁰.

Chlorophyll content serves as a critical indicator of plant health and photosynthetic activity. The decline in chlorophyll content with higher atrazine rates suggests a potential interference with the photosynthetic process¹¹. While the reduction in chlorophyll content may be associated with the herbicide's impact on weeds, the study highlights the need for a nuanced approach to atrazine application to minimize adverse effects on maize crops.

The positive correlation between atrazine application rates and maize grain yield underscores the herbicide's pivotal role in optimizing crop productivity. Higher atrazine doses effectively reduced weed competition, allowing maize plants to allocate resources more efficiently toward grain production.

The increase in biomass and straw yield with higher atrazine rates indicates the herbicide's comprehensive influence on overall crop development¹². Enhanced biomass reflects not only improved grain yield but also increased vegetative plant components, contributing to the resilience and vitality of the maize crop.

The study identifies an optimal range for atrazine application rates, balancing effective weed control with considerations of economic efficiency and environmental impact. The dosage of 1.5

ml/L emerged as the point of maximum efficacy, achieving substantial weed suppression without compromising maize health.

CONCLUSION

The optimization of atrazine application rates represents a delicate dance between weed control efficacy and potential impacts on maize crops. This study contributes valuable insights to the ongoing discourse on responsible herbicide use, urging a nuanced and context-specific approach in the pursuit of sustainable and efficient maize cultivation practices. As agriculture evolves, precision-based weed management strategies become paramount, and these findings contribute to the collective knowledge guiding farmers, agronomists, and policymakers toward more informed decision-making in herbicide application for maize crops.

REFERENCES:

- Hamill AS, Zhang J. Rate and time of bentazon/atrazine application for broadleaf weed control in corn (*Zea mays*). *Weed Tech.* 1997;11(3):549-55.
- Chikoye D, Udensi UE, Lum F. Performance of a new formulation of atrazine for weed control in maize in Nigeria. *J. Food Agric. Environ.* 2006;4(3/4):114.
- Walker SR, Robinson GR, Hargreaves PA. Weed control with atrazine and chlorsulfuron is determined by herbicide availability and persistence in soils. *Aus. J. Agric. Res.* 1997;48(7):1003-10.
- Arslan ZF, Williams MM, Becker R, Fritz VA, Peachey RE, Rabaey TL. Alternatives to atrazine for weed management in processing sweet corn. *Weed Sci.* 2016;64(3):531-9.
- Osipitan OA, Scott JE, Knezevic SZ. Tolpyralate applied alone and with atrazine for weed control in corn. *The J. Agric. Sci.* 2018;10(10):32-9.
- Armel GR, Wilson HP, Richardson RJ, Hines TE. Mesotrione, acetochlor, and atrazine for weed management in corn (*Zea mays*). *Weed Tech.* 2003;17(2):284-90.
- Grichar WJ, Besler BA, Brewer KD. Weed control and grain sorghum (*Sorghum bicolor*) response to postemergence applications of atrazine, pendimethalin, and trifluralin. *Weed Tech.* 2005;19(4):999-1003.
- Eckert RE, Asher JE, Christensen MD, Evans RA. Evaluation of the atrazine-fallow technique for weed control and seedling establishment. *Rangeland Ecol. Manage./J. Range Manage. Arch.* 1974;27(4):288-92.
- Williams MM, Boerboom CM, Rabaey TL. Significance of atrazine in sweet corn weed management systems. *Weed Tech.* 2010;24(2):139-42.
- Krutz LJ, Zablotowicz RM, Reddy KN, Koger Iii CH, Weaver MA. Enhanced degradation of atrazine under field conditions correlates with a loss of weed control in the glasshouse. *Pest Manage. Sci. Form. Pestic. Sci.* 2007;63(1):23-31.
- Chikoye D, Udensi UE, Lum AF. Evaluation of a new formulation of atrazine and metolachlor mixture for weed control in maize in Nigeria. *Crop Protec.* 2005;24(11):1016-20.
- Balyan RS, Kumar S, Malik RK, Panwar RS. Post-emergence efficacy of atrazine in controlling weeds in pearl-millet. *Ind. J. Weed Sci.* 1993;25(1and2):7-11.